

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/124417/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Watson, Sophie E., Hauffe, Heidi C., Bull, Matthew J., Atwood, Todd C., McKinney, Melissa A., Pindo, Massimo and Perkins, Sarah E. ORCID: <https://orcid.org/0000-0002-7457-2699> 2019. Global change-driven use of onshore habitat impacts polar bear faecal microbiota. ISME Journal 13 , pp. 2916-2926. 10.1038/s41396-019-0480-2 file

Publishers page: <http://dx.doi.org/10.1038/s41396-019-0480-2>
<<http://dx.doi.org/10.1038/s41396-019-0480-2>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Global change-driven use of onshore habitat impacts polar bear faecal microbiota

Sophie E. Watson^{1,2}, Heidi C. Hauffe², Matthew J. Bull^{1,2}, Todd C. Atwood³, Melissa A. McKinney⁴, Massimo Pindo⁵ & Sarah E. Perkins^{1,2}

¹*School of Biosciences, Cardiff University, The Sir Martin Evans Building, Museum Avenue, Cardiff, UK*

²*Department of Biodiversity and Molecular Ecology, Research and Innovation Centre, Fondazione Edmund Mach, S. Michele all' Adige (TN), Italy*

³*United States Geological Survey (USGS), University Drive, Anchorage, USA*

⁴*Department of Natural Resource Sciences, McGill University, Ste-Anne-de-Bellevue, Quebec, Canada*

⁵*Department of Genomics and Biology of Fruit Crops, Research and Innovation Centre, Fondazione Edmund Mach, S. Michele all' Adige (TN), Italy*

Corresponding author

Sophie Watson (email: WatsonS2@cardiff.ac.uk, telephone: 02920 875073)

Competing interests

We declare that the authors have no conflicts of interest that might be perceived to influence the results and/or discussion that are reported in this article.

Running title

Land use alters polar bear faecal microbiota

34 **Abstract**

35 The gut microbiota plays a critical role in host health, yet remains poorly studied in wild species. Polar
36 bears (*Ursus maritimus*), key indicators of Arctic ecosystem health and environmental change, are
37 currently affected by rapid shifts in habitat that may alter gut homeostasis. Declining sea ice has led to
38 a divide in the southern Beaufort Sea polar bear subpopulation such that an increasing proportion of
39 individuals now inhabit onshore coastal regions during the open-water period ('onshore bears') while
40 others continue to exhibit their typical behaviour of remaining on the ice ('offshore bears'). We
41 propose that bears that have altered their habitat selection in response to climate change will exhibit a
42 distinct gut microbiota diversity and composition, which may ultimately have important consequences
43 for their health. Here, we perform the first assessment of abundance and diversity in the faecal
44 microbiota of wild polar bears using 16S rRNA Illumina technology. We find that bacterial diversity
45 is significantly higher in onshore bears compared to offshore bears. The most enriched OTU abundance
46 in onshore bears belonged to the phylum Proteobacteria, while the most depleted OTU abundance
47 within onshore bears was seen in the phylum Firmicutes. We conclude that climate-driven changes in
48 polar bear land use are associated with distinct microbial communities. In doing so, we present the first
49 case of global change mediated alterations in the gut microbiota of a free-roaming wild animal.

50

51

52 **Introduction**

53 As an apex predator with vulnerable conservation status [1], the polar bear (*Ursus maritimus*) is widely
54 acknowledged as a key indicator of Arctic ecosystem health [2], a model species for studying the
55 effects of climatic and other anthropogenic stressors in the Arctic [3–5], and a flagship for
56 environmental change [6]. As one of the most ice dependent Arctic marine mammals [7], polar bears
57 require sea ice for long-distance movements, mating and accessing prey [8]. One subpopulation of
58 polar bear, the southern Beaufort Sea subpopulation, is exhibiting a distinct behavioural response to
59 climate-driven changes in sea ice conditions. Historically, these polar bears remained year-round on
60 the sea ice (hereafter referred to as 'offshore bears'), taking advantage of the biologically-productive
61 continental shelf [9]. Since the 2000s, however, substantial declines in the spatial and temporal
62 availability of sea ice in summer and fall [10, 11], extending well beyond the continental shelf, have
63 driven a divide in polar bear behaviour whereby some continue to select the retreating ice habitat
64 ('offshore bears') while others instead adopt a novel behaviour and move to coastal onshore habitat
65 during the reduced ice period ('onshore bears')[12]. The entire subpopulation uses the sea ice during

66 the remainder of the year. Onshore bears have been associated with a range of dietary items that
67 offshore bears are unable to access, notably ‘bone piles’, the remains of locally-harvested bowhead
68 whales (*Balaena mysticetus*), along with the carcasses of fish, birds and caribou (*Rangifer tarandus*)
69 [13]. Conversely, offshore bears primarily consume a traditional diet of ringed seal (*Pusa hispida*),
70 bearded seal (*Erignathus barbatus*) and occasionally beluga whale (*Delphinapterus leucas*) [13],
71 which are inaccessible to onshore bears.

72 Changes in trophic interactions alter the exposure of polar bears to contaminants and novel parasites
73 [14, 15]. For example, ringed seals (available only to offshore bears) are considered to occupy a high
74 trophic position and so typically bioaccumulate higher levels of contaminants than species lower in
75 the trophic chain such as the filter feeders (i.e. bowhead whales) and herbivores (i.e. caribou) [16–18],
76 which are available only to onshore bears. In addition, bone piles, foraged on by onshore bears, are
77 utilised as a food resource by other terrestrial species [13, 19] and lie within comparatively close range
78 of human settlements, such as Kaktovik (70.13° N, 143.62° W) and Deadhorse (70.20° N, 148.46° W).
79 Thus, onshore bears are potentially exposed to (and therefore at greater risk of infection from) novel
80 parasites carried by terrestrial species, including humans and their domestic pets. For example, Atwood
81 *et al.* (2017) [5] found that southern Beaufort Sea polar bears exhibiting onshore behaviour have a
82 greater risk of exposure to *Toxoplasma gondii* and lower exposure to certain contaminants than
83 offshore bears. Thus, onshore bears are exposed to different biotic stressors compared to offshore bears
84 [5, 20], which have the potential to drive variation in the gut microbiota. In humans and mice, for
85 example, helminth infection is associated with significant differences in the community composition
86 of gut bacterial communities [21–23], while contaminants such as herbicides and pesticides have been
87 shown to inhibit the growth of a variety of beneficial gut bacteria [24] and even cause dysbiosis [25].

88 The gut microbiota, a diverse community of bacteria that resides within the gastrointestinal tract, has
89 a long co-evolutionary association with its host [26], carrying out vital nutritional and physiological
90 roles [26–28]. In effect, the regular intestinal development and function of an individual is attributed
91 to an array of specific bacterial groups or species, the composition and diversity of which are a function
92 of complex interactions between host and environment [29]. Despite the importance of the gut
93 microbiota to health, little is understood of the composition or community structure of the gut
94 microbiota of wild fauna [30]. In brown bears (*U. arctos*) however, we know a distinct gut microbiota
95 profile is associated with active bears compared to those in hibernation phase – this specific community
96 of bacteria is thought to play a role promoting adiposity while still maintaining normal gut metabolism
97 [31]. A paucity of knowledge on wild microbiota is particularly concerning considering that in the face
98 of rapid climate change tight host-gut microbiota associations could quickly become decoupled,

99 negating millions of years of co-evolutionary adaptation [26], and yet this too remains poorly
100 understood.

101 A number of studies provide support for an association between host microbial communities and
102 environmental fluctuations. Cold acclimated laboratory mice, for example, harbour a dramatically
103 different gut microbiota composition to those raised at higher temperatures [32], while experimentally
104 induced temperature increases of 2–3 °C cause a 34% loss of microbiota diversity in the common lizard
105 (*Zootoca vivipara*)[33]. Outside a laboratory setting, variations in weather events have been linked to
106 the increased occurrence of gastrointestinal illness in residents of Nunatsiavut, Canada [34]. To the
107 best of our knowledge, however, no study has demonstrated a climate change mediated alteration in
108 the gut microbiota of free-roaming wildlife.

109 The gut microbiota has been examined once before in wild polar bears, specifically those from the
110 Svalbard archipelago belonging to the Barents Sea subpopulation [35]. The authors found a low
111 bacterial diversity, dissimilar to that reported in other Arctic carnivores [36] and wild ursids [31, 37,
112 38], possibly attributed to the methodologies employed (having used 16S rRNA clone libraries as
113 opposed to next generation sequencing techniques) and small sample size [35, 39]. Here we use high-
114 throughput sequencing techniques to conduct the first detailed investigation of the gut microbiota
115 composition of a large sample ($n=112$) of wild southern Beaufort Sea polar bears and to establish the
116 diversity, abundance, and composition of gut bacteria associated with on- and offshore bears. In doing
117 so, we are able to evaluate the effect of a climate driven change in habitat use on microbial
118 composition. Reflecting methods widely used in other gut microbiota studies [40], we use faeces as a
119 proxy of gut microbiota, herein referred to as the faecal microbiota.

120 **Materials and methods**

121 **Polar bear capture and sampling**

122 Polar bears were captured under the United States Geological Survey (USGS) Polar Bear Research
123 Program (Marine Mammal Permit MA690038 to T.C.A.) in an area ranging approximately from
124 Utqiagvik, Alaska (156°W) in the west to Demarcation Point (140°W) at the US-Canada border in the
125 east, and extending from the shoreline to approximately 135 km north on sea ice (with the exception
126 of one individual; Figure 1). In the spring and fall of 2008 and 2009, and the spring of 2010 and 2013,
127 polar bears were encountered via helicopter and immobilized with a remote injection of zolazepam-
128 tiletamine (Telazol®, Fort Dodge Animal Health, Fort Dodge, Iowa, USA, and Warner-Lambert Co.,
129 Groton, Connecticut, USA). A single faecal sample was collected directly from the rectum of each

130 polar bear using a sterile latex glove and immediately transferred to a sterile Whirl-pak bag (Nasco,
131 Fort Atkinson, Wisconsin, USA) for storage. In total, samples were taken from 112 individuals,
132 including 89 adults and 23 subadults, (51 males and 61 females). All samples were stored at -20°C for
133 the duration of the field season (approx. 5 weeks) before being stored at -80°C at the US Geological
134 Survey, Alaska Science Center (Anchorage, Alaska, USA), and subsequently shipped on dry ice to the
135 Fondazione Edmund Mach, Italy (CITES permit IT/IM/2015/MCE/01862 to S.W.).

136 Age of subadults and adults was estimated by extracting and analysing the cementum annuli of a
137 vestigial premolar tooth [41]. In total, 85 of the 112 bears were known to be either onshore or offshore
138 (onshore $n = 46$; offshore $n = 39$; Supplementary Table 1). Individuals were categorised as either
139 'onshore bears' or 'offshore bears' as described in [5]. Briefly, location data collected from satellite
140 collars were used to identify adult females that used land ('onshore') or sea ice ('offshore') in summer
141 and fall [42]. We classified both male and female individuals as onshore bears if they were detected
142 (via genetic identification and cross-referencing with our database of known bears) at hair-snags
143 erected in the fall around bowhead whale bone piles and from biopsy-darting during fall coastal surveys
144 from 2010-2013. An individual was classified as onshore or offshore if spatial or genetic data
145 suggested that the individual was onshore or offshore in summer and/or in the year of capture (for fall-
146 captured bears) or immediately prior to capture (for spring-captured bears). Body condition for each
147 polar bear was estimated using a 'Body Condition Index' metric [43] and was classified as either above
148 or below the mean body condition for our sample set. Year and season of capture was also recorded.

149 **Extraction of bacterial DNA**

150 All faecal matter was collected from inside each sample glove using a sterile cotton swab (APTACA
151 sterile transport swabs, Brescia, Italy). The swab was subsequently vortexed for 10 min in 1ml
152 phosphate-buffered saline solution (PBS) and pelleted by centrifugation at 16 000 g for 12 min. Lysis
153 buffer, 80 μ l, (200 mM NaCl, 100 mM Tris, 20 mM EDTA, 20 mg/ml Lysozyme, pH 8.0); 5 mm
154 stainless steel beads (Qiagen) were added to each sample before a three-minute homogenization step
155 at 30Hz using a Mixer Mill MM200 (Retsch GmbH, Haan, Germany). Samples were then shaken at
156 37°C for 40 minutes Grant-Bio PCMT Thermoshaker (500rpm). Microbial DNA was extracted using
157 the QIAamp® DNA Mini Kits (QIAGEN®, Milan, Italy), following the manufacturer's Buccal Swab
158 Spin Protocol for cotton swabs (QIAamp® DNA Mini and Blood Mini Handbook), but starting from
159 step 2 (addition of Proteinase K).

160

161 **16s rRNA gene amplification and sequencing**

162 Using the bacteria-specific primer set 341F (5' CCTACGGGNGGCWGCAG 3') and 805Rmod (5'
163 GACTACNVGGGTWTCTAATCC 3') (based on Klindworth *et al.* 2013 [44] with degenerate bases)
164 with overhanging Illumina adapters, a ~460 base pair (bp) fragment of the 16S rRNA gene (variable
165 region V3-V4)[45] was amplified using a GeneAmp PCR System 9700 (Thermo Fisher Scientific) and
166 the following steps: 94°C for 5 minutes (one cycle), 95°C for 30 seconds, 55°C for 30 seconds, 72°C
167 for 30 seconds (30 cycles), 72°C for 5 minutes (1 cycle). The PCR products were visualised on a 1.5%
168 agarose gel and purified using Agencourt AMPure XP SPRI beads (Beckman Coulter, Brea, CA, USA)
169 following manufacturer's instructions. Subsequently, Illumina® Nextera XT indices and sequencing
170 adapters (Illumina®) were incorporated using seven cycles of PCR (16S Metagenomic Sequencing
171 Library Preparation, Illumina®). The final libraries were quantified using the Quant-IT PicoGreen
172 dsDNA assay kit (Thermo Fisher Scientific) by the Synergy2 microplate reader (Biotek), pooled in
173 equimolar concentration before sequencing on an Illumina® MiSeq (2x300 bp reads) at the Next
174 Generation Sequencing Platform, Fondazione Edmund Mach in collaboration with the Core Facility,
175 CIBIO, University of Trento, Italy. All samples were sequenced in one Illumina MiSeq Standard Flow
176 Cell targeting a depth of 20 000 reads per sample.

177 **Bioinformatic processing of 16s data**

178 Reads were processed with MICCA v1.5.0 [46]. Briefly, paired-end reads were merged, and pairs
179 diverging by more than 8 bp or overlapping by less than 100 bp were discarded. PCR amplification
180 primers were trimmed (sequences not containing both PCR primer sequences were discarded). Finally,
181 sequences were quality filtered at 0.5 % Expected Error (EE); those displaying greater than 0.5% EE
182 were discarded along with those shorter than 400 bp or containing unknown base calls (N). Using the
183 VSEARCH cluster_smallmem algorithm [47], OTUs were created *de novo* by clustering sequences
184 with 97% sequence identity, discarding chimeric sequences. Taxonomic assignments of representative
185 sequences from each OTU were performed using the RDP Classifier v2.12 in conjunction with RDP
186 16S rRNA training set 15 [48]. OTU sequences were aligned and phylogenetic analysis was performed
187 using Nearest Alignment Space Termination (NAST) and a phylogeny reconstructed using FastTree
188 [49], both via MICCA [46]. The raw sequencing data can be found at the National Centre for
189 Biotechnology Information (NCBI) Sequence Read Archive (SRA) [Accession number:
190 PRJNA542176].

191

192 **Statistical analyses**

193 Following initial processing, singletons were removed and all samples with fewer than 5000 reads
194 were removed using the R package ‘phyloseq’ [50], leaving a total of 511 952 reads across 112
195 samples. The data were rarefied to an equal depth within 90% of the minimum observed sample size
196 (specifically 4571 reads per sample). Generalized Linear Models (GLMs) with a Gamma error function
197 were used to investigate whether metadata (onshore/offshore, age class, sex, body condition, year of
198 capture and season of capture) were associated with alpha diversity of the faecal microbiota (Shannon,
199 Inverse Simpson and Faith’s Phylogenetic Diversity Indices). For Shannon and Faith’s Phylogenetic
200 Diversity measures, an identity link function was used, while a log link function was used when
201 analysing an Inverse Simpson measure of diversity. All multivariate analyses on faecal microbiota
202 structure according to host metadata (on-/offshore, age class, sex, body condition, year of capture and
203 season) were assessed using PERMANOVA, based on Bray-Curtis dissimilarity and weighted
204 UniFrac indices, using the ‘adonis’ function in the R package ‘vegan’ [51]. An important assumption
205 for PERMANOVA is homogenous dispersion of data among groups; for this reason, the ‘betadisper’
206 function in ‘vegan’ was implemented to investigate the homogeneity of data. Data rows containing
207 missing values (NAs) were removed from the dataset prior to conducting the PERMANOVA to ensure
208 matrices were even between variables. To determine the differential abundance of OTUs between on-
209 and offshore bears, sex and season were examined using the R package ‘DESeq2’ [52]. To assess
210 whether the microbiota profiles of polar bears is related to their geographic distribution, a GPS based
211 pairwise distance matrix was constructed using the R package ‘geosphere’ [53] and compared to a
212 PCoA matrix (using both Bray-Curtis and weighted UniFrac) via a Mantel Test. All analyses were
213 carried out using R statistical software package, version 3.2.0 [54]. Data was visualised using the R
214 packages ‘ggplot2’ [55] and ‘metacoder’ [56].

215 **Results**

216 **Faecal microbiota composition**

217 The faecal microbiota of all 112 bears was composed of 1221 operational taxonomic units (OTUs)
218 encompassing 25 bacterial phyla, with prevalence and abundance of specific phyla differing among
219 individuals (Figure 2a). Across the population, the most abundant phyla (which composed 91% of the
220 total reads and were present in all individuals) were Firmicutes (45%), Proteobacteria (25%) and
221 Actinobacteria (21%), making up the core microbiota. All other phyla represented <9% of reads each
222 (Figure 2a), and their prevalence among samples varied between 97% (Bacteroidetes) and 1%
223 (Armatimonadetes, Deferribacteres, Lentisphaerae and Synergistetes). From the total number of reads

224 obtained for the most dominant phylum (Firmicutes), 70% belonged to the class Clostridia, and 99%
225 of those were from the order Clostridiales. The dominant orders for the remaining top bacterial phyla
226 were Enterobacteriales (phyla: Proteobacteria) and Actinomycetales (phyla: Actinobacteria) (Figure
227 2b).

228 **Onshore versus offshore microbiota**

229 Using the subset of bears for which we had on- and offshore information ($n = 85$), we found alpha
230 diversity was significantly higher in on- ($n = 46$) compared to offshore ($n = 39$) bears, for Shannon
231 (adjusted R-squared = 0.06, $F_{1,83} = 6.32$, $P = 0.014$; Figure 3a and Supplementary Table 2) and Inverse
232 Simpson (adjusted R-squared = 0.07, $F_{1,83} = 6.09$, $P = 0.016$; Figure 3b and Supplementary Table 2)
233 indices but not for Faith's Phylogenetic Diversity index (Supplementary Table 3). Beta diversity did
234 not differ between on- and offshore bears when using Bray-Curtis (Supplementary Figure 1) but
235 differed significantly between on- and offshore bears when using a weighted UniFrac metric (adjusted
236 R-squared = 0.03, $F_{1,80} = 2.53$, $P = 0.029$; Supplementary Figure 2). Data dispersion did not
237 significantly differ between on- and offshore bears ($P=0.740$).

238

239 The faecal microbiota of onshore bears consisted of 858 OTUs (19 bacterial phyla; 37 classes)
240 compared to 635 OTUs (21 phyla; 35 classes) for offshore bears, of which 386 were shared between
241 on- and offshore polar bears (Figure 4). Of the total number of OTUs found 472 were unique to
242 onshore bears, and a smaller number of OTUs ($n= 249$) were unique to offshore bears. Eleven OTUs
243 (10 Firmicutes; 1 Proteobacteria) were significantly enriched and 6 OTUs (3 Bacteroidetes; 2
244 Firmicutes; 1 Proteobacteria) were significantly reduced in onshore bears (Figure 5; Supplementary
245 Table 4). The majority (73%; $n = 8$) of OTUs that were enriched in onshore bears belonged to the order
246 Clostridiales (Phylum: Firmicutes), although family level assignment varied across OTUs (Figure 5
247 and Supplementary Table 4). OTUs that were significantly decreased in on- compared to offshore bears
248 varied in taxonomic assignment across taxonomic ranks (Supplementary Table 4). The most enriched
249 OTU abundance in onshore bears belonged to the family Moraxellaceae (Phylum: Proteobacteria),
250 with a 6.78 log₂ fold change in abundance ($P<0.001$), while the most depleted OTU abundance within
251 onshore bears was seen in Clostridiaceae 1 (Phylum: Firmicutes) with a -8.04 log₂ fold change in
252 abundance ($P<0.001$; Supplementary Table 4).

253 The gut microbiota composition of individuals was not associated with their geographic proximity to
254 one another ($P=0.56$ and $P=0.17$; Mantel Test using Bray-Curtis and weighted Unifrac respectively).

255

Ecological factors and the microbiota

When using Faith's Phylogenetic Diversity Index, alpha diversity was significantly higher in females compared to males (adjusted R-squared = 0.30, $F_{2,109} = 25.18$, $P = 0.017$), as well as in fall compared to spring captures (adjusted R-squared = 0.30, $F_{2,109} = 25.18$, $P < 0.001$). However, alpha diversity did not differ with sex, season of capture, body condition, year or age class when using either a Shannon or Inverse Simpson index of diversity and no significant difference in alpha diversity was seen with body condition, year, or age class when using Faith's Phylogenetic Diversity. Beta diversity differed significantly with sex (Bray-Curtis; $P = 0.001$; weighted UniFrac $P = 0.006$) although data dispersion was seen to be significantly different between males and females ($P = 0.018$) and so the PERMANOVA should be interpreted with caution. Beta diversity also differed significantly with and season when using Bray-Curtis ($P = 0.005$) but not weighted UniFrac ($P = 0.184$), where beta dispersion was $P = 0.113$. No differences in beta diversity were seen with year, age class or body condition when using either Bray-Curtis or a weighted UniFrac metric. When investigating the differential abundance of OTUs with sex, DESeq analysis showed that 66 OTUs were significantly different between males and females; 9 OTUs were significantly increased in males compared to females (the largest increase, of 5.40 log fold change, belonging to the family Clostridiales Incertae Sedis XI, phylum: Firmicutes) and 57 OTUs were significantly decreased (the largest decrease, of -10.04 log fold change, being seen in the family Flavobacteriaceae, phylum: Bacteroidetes). For season of capture, DESeq analysis revealed that 15 OTUs were significantly different between fall and spring captures; 2 OTUs were increased in spring compared to fall captures (the largest increase, of 3.01 log fold change, belonging to the family Veillonellaceae, phylum: Firmicutes) and 13 OTUs were significantly decreased (the largest decrease, of -7.50 log fold change, being seen in the family Peptostreptococcaceae, phylum: Firmicutes).

Discussion

Investigating factors which may influence the gut microbiota in a sentinel species experiencing rapid environmental change may improve our understanding of the role of the gut microbiota in wildlife health and conservation. Here we have shown that for the southern Beaufort Sea subpopulation of polar bears alpha diversity and bacterial composition are significantly different in the gut of onshore bears compared to those that remain on the sea ice year-round. As such, our study shows for the first time, that global change driven alterations in habitat use are associated with changes in the gut microbial composition and diversity of a free-ranging species.

287 We detected 25 bacterial phyla, as opposed to just the one (Firmicutes) previously found by Glad *et*
288 *al.* (2010) [35] in wild Barents Sea polar bears. This diversity closely mirrors that seen in other studies
289 utilizing next generation sequencing methods to investigate the gut microbiota of ursids; for example,
290 24 bacterial phyla were detected in wild brown bears [31]. The most abundant phyla in polar bear
291 faeces (Firmicutes, Proteobacteria and Actinobacteria), coincided with those of the core mammalian
292 gut microbiota [26], including that of Asiatic black bears (*Ursus thibetanus*) [38]. Our finding that
293 Firmicutes constituted the majority of OTUs is noteworthy in that increased Firmicutes in genetically
294 obese mice and humans suggests that this phylum plays an important role in promoting adiposity or
295 energy resorption [57], although conflicting studies show no link between Firmicutes levels and
296 obesity/high-fat intake [58]. Interestingly, brown bears gaining weight for hibernation during summer
297 months show simultaneously elevated levels of Firmicutes in the gut [31], implying this phylum may
298 also play a role in synthesising high energy inputs in large carnivores. More specifically, we show that
299 70% of reads assigned to the phylum Firmicutes belonged to the class Clostridia, and subsequently
300 99% were from the order Clostridiales – an outcome that coincides with the results of Glad *et al.*
301 (2010), who showed all except one of the gene clones generated within their study were affiliated with
302 the order Clostridiales. In a study using both wild type and laboratory mice, Hilderbrant *et al.* (2009)
303 [59] showed that levels of Clostridiales greatly increase after prolonged durations of time feeding on
304 a high-fat diet.

305 Within this study we found that alpha diversity of bacterial OTUs was significantly higher in the faecal
306 microbiota of onshore compared to offshore bears when using a Shannon or Inverse Simpson measure,
307 but no association was found between alpha diversity and host metadata (age class, sex, body
308 condition, year or season of capture) when using these indices. Much microbiota work focusing on
309 humans has found sex and age influences microbiota dynamics [60–62]. Although the majority of
310 microbiota research has focused on humans, microbial studies of wild animals are increasing [30] and
311 in some cases wild animals have been shown to follow similar trait-related stratification in microbiota.
312 For example, the presence/absence of specific bacterial taxa were seen to correlate with specific age
313 classes within the gut microbiota of wild ring-tailed lemurs (*Lemur catta*) [63]. Similarly, sex-specific
314 differences in bacterial diversity have been found in, for example, wild rufous mouse lemurs
315 (*Microcebus rufus*), whereby females demonstrated higher bacterial diversity compared to their male
316 counterparts [64]. Further to this, season of capture has been seen to influence the gut microbiota
317 composition. Sommer *et al.* (2016) [31], for example, demonstrated that gut microbial composition of
318 free-roaming brown bears is seasonally altered between summer and winter. This change in bacterial
319 composition is thought to, in part, be influenced by extreme dietary shifts within brown bears between

active and hibernation phase [30]. We also see this seasonal shift in gut microbial composition in other wild animal models such as wild wood mice (*Apodemus sylvaticus*) [65], wild black howler monkey (*Alouatta pigra*) [66], and the giant panda (*Ailuropoda melanoleuca*) [37], probably also attributable to season-driven shifts in diet. None of these factors, however, were found to influence the gut microbiota composition of the polar bears sampled within this study when using a Shannon and Inverse Simpson index of diversity. However, when using Faith's Phylogenetic Diversity (i.e. a metric that characterises only the relatedness or distinctness of species and works under the assumption that different species make unequal contributions to diversity [67]) we see a significant difference in diversity with sex and season only, whereby females had a higher bacterial diversity than males, and fall captures had a higher bacterial diversity than spring captures. Faith's phylogenetic diversity index does not incorporate the relative abundances of taxa within communities, but rather calculates phylogenetic diversity based on the presence or absence of species [68, 69]. Our results therefore imply that for sex and season, there was no difference in alpha diversity when considering the richness and evenness of species, but that there may be a number of species with deep and/or distinct branching that are making an unequal contribution to the diversity of those communities.

We posit that the differences in gut microbiota composition between on- and offshore bears is most likely driven by environmental factors, such as diet, contaminants and parasites which are known to differ between the two groups [70–73] – although this hypothesis is yet to be tested. Diet, as one of the biggest drivers in gut microbial changes [74–76], likely plays the largest role in the observed differences in bacterial diversity. Historically, southern Beaufort Sea polar bears remained offshore hunting ringed seal (*Pusa hispida*) and, to a lesser extent, bearded seal (*Erignathus barbatus*) [77], primarily consuming high-calorie blubber with a specific, restricted nutritional input [78]. In contrast, onshore bears have access to a more varied but less natural diet, including bowhead whale bone piles, which can consist of whale blubber, meat, and viscera, as well the carcasses of fish, birds and caribou (*Rangifer tarandus*) [42, 79, 80], a more varied food source in terms of both species and tissue types.

Not only do onshore bears consume a larger range of food items, but they also likely come into contact with more terrestrial species and their associated bacteria and pathogens. Whale bone piles are utilised by a range of other nearshore/terrestrial scavengers [5, 19] providing an inter-specific focal point for many species with which polar bears do not typically interact. Beach-cast bowhead whale remains frequently lie in close proximity to settlements and towns, increasing the potential for microbiota and pathogen spillover to polar bears from humans, and domestic animals. The high gut microbiota diversity seen in onshore bears may therefore be associated with this complex network of interspecific contacts. A secondary consequence of high inter-species contact could be a higher parasite load and/or

diversity in polar bears, which is associated with high gut microbiota diversity in other species [23, 29, 81].

Understanding the ways in which polar bears respond to climate-change mediated displacement from primary habitat is crucial in discerning their ability to cope with an increasingly changeable and uncertain environment [42]. Future management plans for polar bears could therefore benefit from a better understanding of the relationship between habitat availability, microbiota and health. Our results suggest that climate driven changes in land use by bears leads to changes in gut community composition, but further analyses are needed to determine whether these changes are linked to underlying causes such as diet, parasites and health. It has been suggested that researchers should incorporate health assessments into wildlife conservation practices [82, 83] and long term faecal microbiota monitoring could provide this framework.

Acknowledgments

The authors would like to thank G. Durner, A. Pagano, K. Simac, L. Peacock, and T. Donnelly for capturing and sampling of polar bears, which was funded by the USGS. We are grateful to F. Albonico and the staff at the Conservation Genetics laboratory at the Fondazione Edmund Mach, Italy for their training and guidance. Laboratory facilities and funding for the metataxonomic analyses were provided by the Fondazione E. Mach. This paper was reviewed and approved by USGS under their Fundamental Science Practices policy (<http://www.usgs.gov/fsp>). Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the US Government. S.E.W. is supported by a NERC GW4+ Doctoral Training Partnership studentship from the Natural Environment Research Council [NE/L002434/1].

Contributions

S.E.W.: study concept and design, and acquisition, analysis and interpretation of data, drafting of figures, drafting of manuscript, writing of the manuscript; H.C.H.: study and laboratory supervision, writing of the manuscript and critical revision of the manuscript, obtained funding; M.J.B.: bioinformatic analysis and interpretation of data; T.C.A.: study design, field work and sample collection, data analysis, editing of manuscript, critical revision of the manuscript, obtained funding; M.A.M.: study design, editing of manuscript, critical revision of the manuscript; M.P.: study design and technical support; S.E.P.: study supervision, study concept and design, writing of manuscript and critical revision of the manuscript, obtained funding.

384 **Competing interests**

385 The authors declare no competing interests.

386 **Corresponding author**

387 Correspondence to Sophie Watson (WatsonS2@cardiff.ac.uk)

388 **Supplementary information**

389 Supplementary information is available at ISME's website.

390

391 **References**

- 392 1. Regehr E V, Laidre KL, Akçakaya HR, Amstrup SC, Atwood TC, Lunn NJ, et al.
393 Conservation status of polar bears (*Ursus maritimus*) in relation to projected sea-ice declines.
394 *Biol Lett* 2016; **12**: 20160556.
- 395 2. Amstrup SC, Deweaver ET, Douglas DC, Marcot BG, Durner GM, Bitz CM, et al.
396 Greenhouse gas mitigation can reduce sea-ice loss and increase polar bear persistence. *Nature*
397 2010; **468**: 955–8.
- 398 3. Parmesan C. Ecological and Evolutionary Responses to Recent Climate Change. *Annu Rev*
399 *Ecol Evol Syst* 2006; **37**: 637–669.
- 400 4. McKinney MA, Letcher RJ, Aars J, Born EW, Branigan M, Dietz R, et al. Flame retardants
401 and legacy contaminants in polar bears from Alaska, Canada, East Greenland and Svalbard,
402 2005–2008. *Environ Int* 2011; **37**: 365–74.
- 403 5. Atwood TC, Duncan C, Patyk KA, Nol P, Rhyan J, McCollum M, et al. Environmental and
404 behavioral changes may influence the exposure of an Arctic apex predator to pathogens and
405 contaminants. *Sci Rep* 2017; **7**: 13193.
- 406 6. Derocher AE, Aars J, Amstrup SC, Cutting A, Lunn NJ, Molnár PK, et al. Rapid ecosystem
407 change and polar bear conservation. *Conserv Lett* 2013; **6**: n/a–n/a.
- 408 7. Laidre KL, Stirling I, Lowry LF, Wiig Ø, Heide-Jørgensen MP, Ferguson SH. Quantifying the
409 sensitivity of arctic marine mammals to climate-induced habitat change. *Ecol Appl* 2008; **18**:
410 S97–S125.
- 411 8. Regehr E V., Hunter CM, Caswell H, Amstrup SC, Stirling I. Survival and breeding of polar
412 bears in the southern Beaufort Sea in relation to sea ice. *J Anim Ecol* 2010; **79**: 117–127.
- 413 9. Amstrup SC, Marcot BG, Douglas DC. A bayesian network modeling approach to forecasting
414 the 21st century worldwide status of polar bears. *Arctic Sea Ice Decline: Observations, Projections, Mechanisms, and Implications. Geophysical Monograph 180*. 2008.
415 American Geophysical Union (AGU), pp 213–268.
- 416 10. Stroeve JC, Markus T, Boisvert L, Miller J, Barrett A. Changes in Arctic melt season and
417 implications for sea ice loss. *Geophys Res Lett* 2014; **41**: 1216–1225.
- 418 11. Stern HL, Laidre KL. Sea-ice indicators of polar bear habitat. *Cryosphere* 2016; **10**: 2027–
419 2041.
- 420 12. Schliebe S, Rode KD, Gleason JS, Wilder J, Proffitt K, Evans TJ, et al. Effects of sea ice
421 extent and food availability on spatial and temporal distribution of polar bears during the fall
422 open-water period in the Southern Beaufort Sea. *Polar Biol* 2008; **31**: 999–1010.
- 423 13. Herreman J, Peacock E. Polar bear use of a persistent food subsidy: Insights from non-
424 invasive genetic sampling in Alaska. *Ursus* 2013; **24**: 148–163.
- 425

- 426 14. McKinney MA, Peacock E, Letcher RJ. Sea Ice-associated Diet Change Increases the Levels
427 of Chlorinated and Brominated Contaminants in Polar Bears. *Environ Sci Technol* 2009; **43**:
428 4334–4339.
- 429 15. McKinney MA, Stirling I, Lunn NJ, Peacock E, Letcher RJ. The role of diet on long-term
430 concentration and pattern trends of brominated and chlorinated contaminants in western
431 Hudson Bay polar bears, 1991–2007. *Sci Total Environ* 2010; **408**: 6210–6222.
- 432 16. Dietz R, Riget F, Cleemann M, Aarkrog A, Johansen P, Hansen JC. Comparison of
433 contaminants from different trophic levels and ecosystems. *Sci Total Environ* 2000; **245**: 221–
434 231.
- 435 17. Hoekstra PF, O’Hara TM, Fisk AT, Borgå K, Solomon KR, Muir DCG. Trophic transfer of
436 persistent organochlorine contaminants (OCs) within an Arctic marine food web from the
437 southern Beaufort–Chukchi Seas. *Environ Pollut* 2003; **124**: 509–522.
- 438 18. Bentzen TW, Follmann EH, Amstrup SC, York GS, Wooller MJ, Muir DCG, et al. Dietary
439 biomagnification of organochlorine contaminants in Alaskan polar bears. *Can J Zool* 2008;
440 **86**: 177–191.
- 441 19. Miller S, Wilder J, Wilson RR. Polar bear–grizzly bear interactions during the autumn open-
442 water period in Alaska. *J Mammal* 2015; gyl140.
- 443 20. McKinney MA, Atwood TC, Pedro S, Peacock E. Ecological change drives a decline in
444 mercury concentrations in southern Beaufort Sea polar bears. *Environ Sci Technol* 2017; **51**:
445 7814–7822.
- 446 21. Walk ST, Blum AM, Ewing SA-S, Weinstock J V, Young VB. Alteration of the murine gut
447 microbiota during infection with the parasitic helminth *Heligmosomoides polygyrus*. *Inflamm*
448 *Bowel Dis* 2010; **16**: 1841–9.
- 449 22. Lee SC, Tang MS, Lim YAL, Choy SH, Kurtz ZD, Cox LM, et al. Helminth Colonization Is
450 Associated with Increased Diversity of the Gut Microbiota. *PLoS Negl Trop Dis* 2014; **8**:
451 e2880.
- 452 23. Kreisinger J, Bastien G, Hauffe HC, Marchesi J, Perkins SE. Interactions between multiple
453 helminths and the gut microbiota in wild rodents. *Philos Trans R Soc Lond B Biol Sci* 2015;
454 **370**: 20140295-.
- 455 24. Shehata AA, Schrödl W, Aldin AA, Hafez HM, Krüger M. The Effect of Glyphosate on
456 Potential Pathogens and Beneficial Members of Poultry Microbiota In Vitro. *Curr Microbiol*
457 2013; **66**: 350–358.
- 458 25. Joly C, Gay-Quéheillard J, Léké A, Chardon K, Delanaud S, Bach V, et al. Impact of chronic
459 exposure to low doses of chlorpyrifos on the intestinal microbiota in the Simulator of the
460 Human Intestinal Microbial Ecosystem (SHIME®) and in the rat. *Environ Sci Pollut Res*
461 2013; **20**: 2726–2734.
- 462 26. Ley RE, Lozupone CA, Hamady M, Knight R, Gordon JI. Worlds within worlds: evolution of
463 the vertebrate gut microbiota. *Nat Rev Microbiol* 2008; **6**: 776–788.
- 464 27. Maynard CL, Elson CO, Hatton RD, Weaver CT. Reciprocal interactions of the intestinal
465 microbiota and immune system. *Nature* 2012; **489**: 231–41.
- 466 28. Xing M, Hou Z, Yuan J, Liu Y, Qu Y, Liu B. Taxonomic and functional metagenomic
467 profiling of gastrointestinal tract microbiome of the farmed adult turbot (*Scophthalmus*
468 *maximus*). *FEMS Microbiol Ecol* 2013; **86**: 432–43.
- 469 29. Round JL, Mazmanian SK. The gut microbiota shapes intestinal immune responses during
470 health and disease. *Nat Rev Immunol* 2009; **9**: 313–23.
- 471 30. Pascoe EL, Hauffe HC, Marchesi JR, Perkins SE. Network analysis of gut microbiota
472 literature: an overview of the research landscape in non-human animal studies. *ISME J* 2017;
473 **11**: 2644–2651.
- 474 31. Sommer F, Ståhlman M, Ilkayeva O, Arnemo JM, Kindberg J, Josefsson J, et al. The Gut
475 Microbiota Modulates Energy Metabolism in the Hibernating Brown Bear *Ursus arctos*. *Cell*

- 476 *Rep* 2016; **14**: 1655–1661.
- 477 32. Chevalier C, Stojanović O, Colin DJ, Suarez-Zamorano N, Tarallo V, Veyrat-Durebex C, et
 478 al. Gut Microbiota Orchestrates Energy Homeostasis during Cold. *Cell* 2015; **163**: 1360–1374.
- 479 33. Bestion E, Jacob S, Zinger L, Di Gesu L, Richard M, White J, et al. Climate warming reduces
 480 gut microbiota diversity in a vertebrate ectotherm. *Nat Ecol Evol* 2017; **1**: 0161.
- 481 34. Harper SL, Edge VL, Schuster-Wallace CJ, Berke O, McEwen SA. Weather, Water Quality
 482 and Infectious Gastrointestinal Illness in Two Inuit Communities in Nunatsiavut, Canada:
 483 Potential Implications for Climate Change. *Ecohealth* 2011; **8**: 93–108.
- 484 35. Glad T, Bernhardsen P, Nielsen KM, Brusetti L, Andersen M, Aars J, et al. Bacterial diversity
 485 in faeces from polar bear (*Ursus maritimus*) in Arctic Svalbard. *BMC Microbiol* 2010; **10**: 10.
- 486 36. Glad T, Kristiansen VF, Nielsen KM, Brusetti L, Wright A-DG, Sundset MA. Ecological
 487 characterisation of the colonic microbiota in arctic and sub-arctic seals. *Microb Ecol* 2010; **60**:
 488 320–30.
- 489 37. Xue Z, Zhang W, Wang L, Hou R, Zhang M, Fei L, et al. The bamboo-eating giant panda
 490 harbors a carnivore-like gut microbiota, with excessive seasonal variations. *MBio* 2015; **6**:
 491 e00022-15.
- 492 38. Song C, Wang B, Tan J, Zhu L, Lou D, Cen X. Comparative analysis of the gut microbiota of
 493 black bears in China using high-throughput sequencing. *Mol Genet Genomics* 2017; **292**: 407–
 494 414.
- 495 39. Mardis ER. The impact of next-generation sequencing technology on genetics. *Trends Genet*
 496 2008; **24**: 133–141.
- 497 40. Thomas V, Clark J, Doré J. Fecal microbiota analysis: an overview of sample collection
 498 methods and sequencing strategies. *Future Microbiol* 2015; **10**: 1485–1504.
- 499 41. Calvert W, Ramsay MA. Evaluation of age determination of polar bears by counts of
 500 cementum growth layer groups. *Ursus* . International Association for Bear Research and
 501 Management. , **10**: 449–453
- 502 42. Atwood TC, Peacock E, McKinney MA, Lillie K, Wilson R, Douglas DC, et al. Rapid
 503 environmental change drives increased land use by an Arctic marine predator. *PLoS One*
 504 2016; **11**: e0155932.
- 505 43. Cattet MR., Caulkett NA, Obbard ME, Stenhouse GB. A body-condition index for ursids. *Can*
 506 *J Zool* 2002; **80**: 1156–1161.
- 507 44. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, et al. Evaluation of general
 508 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based
 509 diversity studies. *Nucleic Acids Res* 2013; **41**: e1.
- 510 45. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-high-
 511 throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME*
 512 *J* 2012; **6**: 1621–4.
- 513 46. Albanese D, Fontana P, De Filippo C, Cavalieri D, Donati C. MICCA: a complete and
 514 accurate software for taxonomic profiling of metagenomic data. *Sci Rep* 2015; **5**: 9743.
- 515 47. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source tool
 516 for metagenomics. *PeerJ* 2016; **4**: e2584.
- 517 48. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of
 518 rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 2007; **73**: 5261–7.
- 519 49. Price MN, Dehal PS, Arkin AP. Fasttree: Computing large minimum evolution trees with
 520 profiles instead of a distance matrix. *Mol Biol Evol* 2009; **26**: 1641–1650.
- 521 50. McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and
 522 Graphics of Microbiome Census Data. *PLoS One* 2013; **8**: e61217.
- 523 51. Dixon P. VEGAN, a package of R functions for community ecology. *J Veg Sci* 2003; **14**: 927–
 524 930.
- 525 52. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-

seq data with DESeq2. *Genome Biol* 2014; **15**: 550.

53. Hijmans RJ, Williams E, Vennes C. Package ‘geosphere’. *Spherical Trigonometry* 2017.

54. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. 2018. Vienna, Austria.

55. Wickham H. ggplot2: Elegant graphics for data analysis. 2016. Springer-Verlag.

56. Foster ZSL, Sharpton TJ, Grünwald NJ. Metacoder: An R package for visualization and manipulation of community taxonomic diversity data. *PLOS Comput Biol* 2017; **13**: e1005404.

57. Ley RE, Turnbaugh PJ, Klein S, Gordon JI. Microbial ecology: human gut microbes associated with obesity. *Nature* 2006; **444**: 1022–3.

58. Fernandes J, Su W, Rahat-Rozenbloom S, Wolever TMS, Comelli EM. Adiposity, gut microbiota and faecal short chain fatty acids are linked in adult humans. *Nutr Diabetes* 2014; **4**: e121–e121.

59. Hildebrandt MA, Hoffmann C, Sherrill–Mix SA, Keilbaugh SA, Hamady M, Chen Y, et al. High-Fat diet determines the composition of the Murine gut microbiome independently of obesity. *Gastroenterology* 2009; **137**: 1716–1724.e2.

60. Mueller S, Saunier K, Hanisch C, Norin E, Alm L, Midtvedt T, et al. Differences in fecal microbiota in different European study populations in relation to age, gender, and country: a cross-sectional study. *Appl Environ Microbiol* 2006; **72**: 1027–33.

61. Koenig JE, Spor A, Scalfone N, Fricker AD, Stombaugh J, Knight R, et al. Succession of microbial consortia in the developing infant gut microbiome. *Proc Natl Acad Sci* 2011; **108**: 4578–4585.

62. Dominianni C, Sinha R, Goedert JJ, Pei Z, Yang L, Hayes RB, et al. Sex, Body Mass Index, and Dietary Fiber Intake Influence the Human Gut Microbiome. *PLoS One* 2015; **10**: e0124599.

63. Bennett G, Malone M, Sauter ML, Cuozzo FP, White B, Nelson KE, et al. Host age, social group, and habitat type influence the gut microbiota of wild ring-tailed lemurs (*Lemur catta*). *Am J Primatol* 2016; **78**: 883–892.

64. Avelo T, Laakkonen J, Jernvall J. Population- and Individual-Level Dynamics of the Intestinal Microbiota of a Small Primate. *Appl Environ Microbiol* 2016; **82**: 3537–45.

65. Maurice CF, CL Knowles S, Ladau J, Pollard KS, Fenton A, Pedersen AB, et al. Marked seasonal variation in the wild mouse gut microbiota. *ISME J* 2015; **9**: 2423–2434.

66. Amato KR, Leigh SR, Kent A, Mackie RI, Yeoman CJ, Stumpf RM, et al. The Gut Microbiota Appears to Compensate for Seasonal Diet Variation in the Wild Black Howler Monkey (*Alouatta pigra*). *Microb Ecol* 2015; **69**: 434–443.

67. Faith DP. Conservation evaluation and phylogenetic diversity. *Biol Conserv* 1992; **61**: 1–10.

68. Cadotte MW, Jonathan Davies T, Regetz J, Kembel SW, Cleland E, Oakley TH. Phylogenetic diversity metrics for ecological communities: integrating species richness, abundance and evolutionary history. *Ecol Lett* 2010; **13**: 96–105.

69. Berg M, Stenuit B, Ho J, Wang A, Parke C, Knight M, et al. Assembly of the *Caenorhabditis elegans* gut microbiota from diverse soil microbial environments. *ISME J* 2016; **10**: 1998–2009.

70. Bentzen TW, Follmann EH, Amstrup SC, York GS, Wooller MJ, O ’hara TM. Variation in winter diet of southern Beaufort Sea polar bears inferred from stable isotope analysis.

71. Schliebe S, Rode KD, Gleason JS, Wilder J, Proffitt K, Evans TJ, et al. Effects of sea ice extent and food availability on spatial and temporal distribution of polar bears during the fall open-water period in the Southern Beaufort Sea. *Polar Biol* 2008; **31**: 999–1010.

72. McKinney MA, Atwood TC, Iverson SJ, Peacock E. Temporal complexity of southern Beaufort Sea polar bear diets during a period of increasing land use. *Ecosphere* 2017; **8**: e01633.

73. Atwood TC, Duncan C, Patyk KA, Nol P, Rhyan J, McCollum M, et al. Environmental and behavioral changes may influence the exposure of an Arctic apex predator to pathogens and contaminants. *Sci Rep* 2017; **7**: 13193.
74. Muegge BD, Kuczynski J, Knights D, Clemente JC, Gonzalez A, Fontana L, et al. Diet Drives Convergence in Gut Microbiome Functions Across Mammalian Phylogeny and Within Humans. *Science* (80-) 2011; **332**: 970–974.
75. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 2013; **505**: 559–563.
76. Carmody RN, Gerber GK, Luevano JM, Gatti DM, Somes L, Svenson KL, et al. Diet Dominates Host Genotype in Shaping the Murine Gut Microbiota. *Cell Host Microbe* 2015; **17**: 72–84.
77. Stirling I, Archibald WR. Aspects of Predation of Seals by Polar Bears. *J Fish Res Board Canada* 1977; **34**: 1126–1129.
78. Rode KD, Robbins CT, Nelson L, Amstrup SC. Can polar bears use terrestrial foods to offset lost ice-based hunting opportunities? *Front Ecol Environ* 2015; **13**: 138–145.
79. Herreman J, Peacock E. Polar bear use of a persistent food subsidy: Insights from non-invasive genetic sampling in Alaska. *Ursus* 2013; **24**: 148–163.
80. McKinney MA, Atwood TC, Iverson SJ, Peacock E. Temporal complexity of southern Beaufort Sea polar bear diets during a period of increasing land use. *Ecosphere* 2017; **8**: e01633.
81. Broadhurst MJ, Ardeshtir A, Kanwar B, Mirpuri J, Gundra UM, Leung JM, et al. Therapeutic helminth infection of macaques with idiopathic chronic diarrhea alters the inflammatory signature and mucosal microbiota of the colon. *PLoS Pathog* 2012; **8**: e1003000.
82. Deem SL, Karesh WB, Weisman W. Putting theory into practice: wildlife health in conservation. *Conserv Biol* 2008; **15**: 1224–1233.
83. Patyk KA, Duncan C, Nol P, Sonne C, Laidre K, Obbard M, et al. Establishing a definition of polar bear (*Ursus maritimus*) health: a guide to research and management activities. *Sci Total Environ* 2015; **514**: 371–8.

Figure legends

Figure 1. Map of study area showing the sampling locations of 112 southern Beaufort Sea polar bears along the north coast of Alaska. Inset map shows the location of the study area, highlighting that one sample originates from a more northerly location than the others.

Figure 2. a) Stacked bar chart of the relative abundance of 25 bacterial phyla in the faecal microbiota of 112 southern Beaufort Sea polar bears. Phyla in the legend are listed in order of decreasing abundance **b)** Inset is a metacoder heatmap plotted to order level: each node moving from the centre outwards represents a different taxonomic rank, whereby kingdom is the centre and nodes representing order appear on the outer edges. The map is weighted and coloured by read abundance.

Figure 3. Violin plots of alpha diversity within the faecal microbiota of 85 southern Beaufort Sea polar bears for which ‘onshore/offshore’ land use is known (see text for definitions): **a)** Shannon diversity index **b)** Inverse Simpson diversity index. Violin plots combine a box plot with a density plot, and as such the width of each plot corresponds to the distribution of the data.

Figure 4. Total number of OTUs in the faecal microbiota of ‘onshore’ and ‘offshore’ bears, by bacterial Class. Inset shows shared OTUs by onshore (green) and offshore (blue) bears.

626 **Figure 5.** Differential OTU abundance of onshore compared to offshore bears from DESeq2 analysis,
627 plotted with individual OTU number and associated family assignment.
628
629